THE RECOVERY KERNEL

How to Navigate the Recovery Kernel

For People Recovering from Caffeine

Coffee, energy drinks, pre-workout, caffeine pills and powders. A step-by-step route: what to read, in what order, and what to expect. Read the note on sources in Section 1 first: unlike the other guides in this series, this one is assembled from cross-references, not from a dedicated Kernel chapter.

> **HARM REDUCTION RESOURCE — NOT MEDICAL ADVICE.** Caffeine is the mildest substance in this series, and it is still a drug. **Get emergency care for chest pain, a pounding or irregular heartbeat that will not settle, fainting, seizure, confusion, or repeated vomiting after a large amount of caffeine.** Powdered and pill forms are the most dangerous, because a small measuring error can be a large overdose.
>
> **The Kernel's specific heart warning:** in a CYP1A2 slow metabolizer, high-dose caffeine, such as very large energy drinks or several in a day, carries an elevated cardiovascular risk. If you have a heart condition, high blood pressure, palpitations, or are pregnant, talk to your clinician before you change anything.
>
> Crisis: call or text 988 · SAMHSA: 1-800-662-4357

1. Before you start

This is a route through the Recovery Kernel for people whose caffeine use is costing more than it gives: anxiety, poor sleep, a racing heart, a cycle of crashing and re-dosing, or caffeine standing in for rest. The Kernel does not treat this as a moral matter. Its frame is neurochemical: caffeine blocks adenosine, the brain's "tired" signal, and the sleep that clears it is when repair happens, so late or heavy caffeine borrows energy against tomorrow's sleep. How it lands depends on two genes it names: COMT (how quickly you clear dopamine and stress chemicals) and CYP1A2 (how quickly you clear caffeine). Six rules follow.

- **Sleep is the thing at stake.** Caffeine's main cost, in the Kernel's account, is the sleep it interferes with, since sleep is when the brain repairs.
- **Know your CYP1A2.** A slow metabolizer keeps caffeine in the body for 7 to 9 hours, so a mid-afternoon drink is still active at midnight.
- **Know your COMT.** For a slow-COMT "Sage," the Kernel says to cut caffeine significantly or eliminate it. For a fast-COMT "Warrior," moderate caffeine is tolerated better, but a cut-off still applies.
- **Energy drinks are the higher-risk form.** Large doses, added stimulants, and a synthetic folic acid blend that does not suit MTHFR variants.
- **Caffeine is not a neutral bridge in recovery from something else.** The Kernel says using it to power through the fatigue of recovery fights the sleep the repair depends on.
- **Do not wait for test results to start.** The genome refines the plan (Main Book, Quick-Start Step 3; Supp. Ch. 5, Step 1b), but the timing rules can start today.

A note on sources (please read). The Kernel files I worked from have no chapter on caffeine. Caffeine appears in scattered places: the Warrior and Sage tables and the substance-avoidance database in Supp. Ch. 6 (Stay Non-Addicted); the Compendium's CYP1A2 passage; Protocol 34 (Cognitive Stabilization); the substitution warning in Supp. Ch. 32 (methamphetamine); the energy note in Supp. Ch. 20 (kratom); and the coffee drinks in Supp. Ch. 49 (Bartender's Friend). Several of these point to an "energy drink intelligence guide" (for the CYP1A2 cut-off, a melatonin timing protocol, and a "strategic caffeine architecture") that is not in the files I have. So this guide has no caffeine withdrawal timeline, product table, or dose ladder from the Kernel, and it does not invent them. Where I add general medical information, it is labeled "added by this guide." If you have the energy drink guide, it should override this one.

How chapters are cited. "Main Book" is the Recovery Kernel Main Book (Second Edition, Centurian). "Compendium" is the Complete Protocol Compendium. "Supp. Ch. N" is chapter N of the Supplemental Book. Section numbers (§) refer to sections inside those chapters.

For caffeine, watch for a second or third dose late in the day to fight an afternoon slump — that slump is often the tail of the first dose, and the new one borrows directly against tonight's sleep.

Redosing is not the only version of this. Watching peers in a treatment setting negotiate a relapse, getting confident advice from someone untrained in your specific situation, or noticing a prescriber treat a prescribed substance more casually than an illicit one can all trigger the same pattern: a felt doubt read as "the plan must be wrong," answered by abandoning it. See **The Commandeering Map**, a companion to this series, for the fuller pattern, including how it can differ by genetic profile and what helps when practice-based resistance alone is not enough.

2. The route at a glance

Steps 0 and 1 come first. Steps 2 to 4 can run at the same time. Steps 5 to 8 overlap and are the work of the first two months. Step 9 is everything after. "P" means Compendium Protocol; "C" means Centurian Paper.

Step	When	What you do	Turn to
0	Day 0	Triage: heart, powders and pills, other stimulants, pregnancy.	Supp. Ch. 6 (avoidance database) Compendium §15.4 (CYP1A2) Main Book, Quick-Start Step 1
1	Days 0–7	Know your pattern: what, how much, and when. What is it doing for you?	Supp. Ch. 6 §4.1 Protocol 34 §34.5
2	Weeks 1–4	Read the Main Book in order.	Main Book: Who This Book Is For, Quick-Start, Part One (Ch. 1–5) Supp. Ch. 2, 3
3	Weeks 1–4	Get the genes that matter: CYP1A2 and COMT first.	Supp. Ch. 5 Step 1 Supp. Ch. 6 §4.3 Compendium P06
4	Weeks 2–6	Resolve your profile.	Supp. Ch. 5 Steps 2–3, 6 Supp. Ch. 6 §4.1–4.3

Step	When	What you do	Turn to
5	Weeks 1–8	Cut back and re-time: the noon cut-off, a ceiling, and no stacking.	Supp. Ch. 6 (avoidance database) Protocol 34 §34.5
6	Weeks 1–8	Repay the sleep debt.	Protocol 34 §34.5 Compendium P17; C-32
7	Weeks 1–12	Build energy and calm without it: movement, meals, water, and the constitution's architecture.	Supp. Ch. 6 §4.1 Supp. Ch. 20 §5 Compendium P12, P19
8	If it applies	If caffeine is bridging another recovery, or a swap for something else, handle it.	Supp. Ch. 32 §3 Supp. Ch. 20 §5–6 Supp. Ch. 6
9	Ongoing, Year 1 to 10	Quarterly check and stewardship.	Kernel quarterly self-check Supp. Ch. 44, 6, 38 Supp. Ch. 5 Step 6

3. The route, step by step

Step 0 — Triage (Day 0, before everything)

Check three things. First, form and amount. The Kernel's example of a risky pattern is very large energy drinks (its example is a 300 mg can) or several in a day, in a CYP1A2 slow metabolizer, and it warns against layering stimulant supplements (caffeine, tyramine-rich foods, yohimbine, synephrine) without knowing your CYP1A2 and COMT status. Second, other stimulants and heart symptoms: if you also use cocaine, methamphetamine, or a prescription stimulant, or you have palpitations, chest pain, or high blood pressure, get medical advice before changing anything and treat the heart symptoms as real. Third, pregnancy or a heart condition: ask your clinician what amount is safe for you.

> **Added by this guide (general medical information, not from the Kernel).** Healthy adults are generally advised that about 400 mg of caffeine a day is not linked to dangerous effects, and pregnancy guidance is lower (commonly about 200 mg). Caffeine powder and concentrated pills are especially risky, because a teaspoon can hold a dangerous amount. These are general figures. Your own limit, especially with a slow CYP1A2 result or a heart condition, may be much lower, and the Kernel's ceiling for a slow metabolizer is lower still (Step 5).

Turn to: Supp. Ch. 6 (the substance-avoidance database, including the CYP1A2 row and the rule about layering stimulant supplements). Compendium Section 15.4 (the secondary metabolic gateways: CYP1A2). Main Book, Quick-Start Boot Sequence, Step 1. Compendium Protocol 01, Emergency Triage, for an acute crisis state.

Step 1 — Know your pattern (Days 0–7)

For a week, write down every source, the amount if you can tell, and the time. Include energy drinks, pre-workout, tea, cola, chocolate, and any pills. Then ask what it was doing for you. The Kernel's lists suggest the usual answers. Energy or motivation: the "understimulated Warrior" (COMT Val/Val, often with MTHFR variants) reaching for stimulants, high-dose caffeine among them, for engagement. Calm through numbness: not typical, since caffeine adds arousal, but the Sage may be using it to power through the fatigue that anxiety and poor sleep produce. Focus: for some people, untreated ADHD. Getting through the day on too little sleep: the most common answer, and the one the Kernel says caffeine hides rather than fixes. Write the answer down, because it decides what replaces it.

Turn to: Supp. Ch. 6 §4.1 (the two constitutions), §4.3 (the full vulnerability matrix). Main Book Ch. 3 (COMT) and Ch. 5 (phenotyping). Compendium Protocol 34 (Cognitive Stabilization) §34.5 (the caffeine reduction row).

Step 2 — Read the Main Book (Weeks 1–4)

Read it in order once you are settled. If you are overwhelmed, the Quick-Start Boot Sequence is designed to be read on its own. Read "Who This Book Is — and Is Not — For" first: it names situations where the supplement protocols need medical modification, including active psychosis, pregnancy, and severe liver disease.

Read in this order: Who This Book Is For → Quick-Start Boot Sequence → Part One: Ch. 1 (Why Willpower Isn't Enough), Ch. 2 (MTHFR), Ch. 3 (COMT; the Warrior and Sage profiles decide most of your caffeine advice), Ch. 4 (CYP450, the drug-metabolism chapter), Ch. 5 (When Systems Collide) → Part Two, the case files (Chapter 6). Ch. 8 (Oxytocin Protocol) is experimental and specialist-only. Ch. 9 (Epigenetic Playground) belongs to Step 9. Appendix A and Appendix B are reference.

Then add: Supp. Ch. 2 (Clarity · Restoration · Manning). The Kernel has no caffeine case study. Supp. Ch. 6 (Stay Non-Addicted), the prevention chapter that carries the caffeine advice. Supp. Ch. 3 (The Addict Virgin Problem) recalibrates timelines. Supp. Ch. 1 (Why They Left You There) is optional reading for the shame.

Step 3 — Get the genes that matter (Weeks 1–4)

The Kernel's recommended route is 30× whole-genome sequencing, which returns a VCF file, a one-time asset you keep. A consumer SNP array is an acceptable start, and a clinical pharmacogenomic panel adds clinical-grade CYP calls. Homocysteine, B12, and RBC folate are functional markers you can recheck quarterly.

For caffeine, two genes lead. **CYP1A2** (the rs762551 variant): the Compendium says slow metabolizers suffer prolonged sympathetic nervous system activation after caffeine, and the Kernel's database gives their half-life as 7 to 9 hours. **COMT** Val158Met: Met/Met (Sage, slow) as the profile most hurt by caffeine, Val/Val (Warrior, fast) as the one that tolerates it better and is more drawn to stimulants. Also relevant: **MTHFR** (energy drinks' B-vitamin blends contain synthetic folic acid, which the Kernel says people with MTHFR variants should avoid in favor of 5-MTHF) and **DRD2** A1 (reward deficiency, which makes any reliable stimulant compelling). Until results arrive, the Kernel's free phenotype clues: severe caffeine-induced anxiety, feeling overwhelmed by stimulation, and a past preference for depressants point to a slow-COMT profile; thriving on deadlines, feeling flat without a challenge, and a past preference for stimulants point to a fast-COMT profile. If you get very jittery or wired from a small amount, or cannot sleep after an afternoon coffee, that is worth telling your clinician.

> **Confirm what you are ordering.** Supp. Ch. 5 specifies 30× whole-genome sequencing with VCF output. "WGBS" more commonly names whole-genome bisulfite sequencing, a DNA-methylation assay that is not built to call variants such as MTHFR C677T. Ask the lab for a standard variant file (VCF).

Turn to: Supp. Ch. 5, Step 1. Supp. Ch. 6 §4.3. Compendium Section 15.4 (CYP1A2), Protocol 06 (Phenotype Assessment). Main Book Ch. 2 and Ch. 3 self-assessments.

Step 4 — Resolve your profile (Weeks 2–6)

Write your profile down and carry key results on a card.

If your result is...	What changes	Source
CYP1A2 slow (the rs762551 "CC" pattern) plus COMT Met/Met	The Compendium's worst case: standard caffeine acts as a severe chemical stressor, worsening anxiety and undermining early recovery. Cut caffeine sharply or eliminate, with no caffeine after noon.	Compendium §15.4; Supp. Ch. 6
CYP1A2 slow alone	Half-life 7–9 hours. High-dose caffeine carries specifically elevated cardiovascular risk. The Kernel's ceiling is 100–150 mg a day, with none after noon.	Supp. Ch. 6 (avoidance database)
COMT Met/Met (Sage)	Caffeine is an amplifier of the Sage's vulnerability: it compounds overstimulation, which the Kernel says can drive depressant-seeking. Reduce significantly or eliminate. Build calm, not stimulation.	Supp. Ch. 6 §4.1, §4.3
COMT Val/Val (Warrior)	Moderate caffeine is tolerated better, but the CYP1A2-based cut-off still applies. Energy drinks and nicotine are named as entry points to harder stimulants, so the intensity has to come from exercise, competition, and mastery.	Supp. Ch. 6 §4.1, §4.3
MTHFR variants	Avoid the synthetic folic acid in energy drinks' B-vitamin blends. Use 5-MTHF. An untreated methylation deficit is part of why the low energy felt bad.	Supp. Ch. 6; Main Book Ch. 2

If your result is...	What changes	Source
DRD2 A1	Reward deficiency: build several genuine reward sources so that a caffeinated one is not the only one.	Supp. Ch. 6 §4.2

Turn to: Supp. Ch. 5, Steps 2, 3, and 6 (gene systems, combined profiles, profile template). Supp. Ch. 6 §4. Main Book Ch. 5 and Appendix A.

Step 5 — Cut back and re-time (Weeks 1–8)

The Kernel's rules, in the order that matters most. No caffeine after noon if you are a CYP1A2 slow metabolizer, since a 3 p.m. energy drink is still active at midnight. A ceiling of 100 to 150 mg a day for a slow metabolizer, and no very large energy drinks or stacking of several. Do not layer stimulant supplements on top. During acute reconstruction after another substance (Protocol 34), reduce or eliminate caffeine, because it raises excitatory activity and lowers GABA function in a system that is already out of balance, and the Kernel rates this as high impact. For a Sage, aim lower still, toward none.

> **Added by this guide (general medical information, not from the Kernel).** Caffeine withdrawal is a recognized condition. Typical features are headache, fatigue, low mood, trouble concentrating, and irritability, starting roughly 12 to 24 hours after the last intake, peaking within a day or two, and usually fading within about a week. Cutting down gradually (for example by shrinking the amount over one to two weeks) usually makes it milder than stopping at once. Stopping is not medically dangerous in the way alcohol or benzodiazepine withdrawal can be, but a bad headache or heart symptoms deserve a clinician's advice.

One tension inside the Kernel to know about. Its Bartender's Friend chapter includes coffee-based drinks (for example a cold brew and cherry drink for the dopamine-deficit profile, and a "Norm" mint coffee drink) and describes cold brew as a slower-release caffeine source for a "rapid metabolizer." That sits awkwardly with the Sage rule above, and with its own advice to cut caffeine during acute reconstruction. Treat the coffee drinks as a choice for a Warrior in stable recovery, not for a slow-COMT or slow-CYP1A2 person.

Turn to: Supp. Ch. 6 (the substance-avoidance database: CYP1A2, the rule on layering stimulants). Compendium Protocol 34 (Cognitive Stabilization) §34.5. Supp. Ch. 49 (Bartender's Friend), with the caution above.

Step 6 — Repay the sleep debt (Weeks 1–8)

The Kernel's position is that sleep is the highest-priority repair intervention, and that stimulants defer the clearance of adenosine and other waste products without removing the need for it: "every hour of stimulant-forced wakefulness adds to the debt," to be collected with interest. Protocol 34 puts seven to nine hours as the most effective single way to reduce intrusive-thought intensity during reconstruction, and it pairs it with regular meals containing protein and fat, since a glucose crash triggers cortisol and anxiety. In practice: a fixed wake time, a real wind-down, and the caffeine cut-off from Step 5. The Kernel's sleep protocol (The Nightly Restoration) and its insomnia paper apply, and it points to a low-dose melatonin timing protocol in the energy drink guide, which I do not have.

Turn to: Compendium Protocol 34 (Cognitive Stabilization) §34.5, Protocol 17 (The Nightly Restoration), Centurian Paper C-32 (The Insomnia Architecture), Protocol 18 (Eating for Restoration). The Kernel's passage on stimulants and sleep debt ("What the Biology Actually Requires").

Step 7 — Build energy and calm without it (Weeks 1–12)

The Kernel's natural replacements for what caffeine was doing. For energy: the Kernel says vigorous exercise, even 20 to 30 minutes, produces the fastest natural dopamine and norepinephrine activation, and a cold shower does too, and that a supplement stack is not an acute energy substitute. For a stable base: regular meals with protein and fat, and hydration, because the Compendium says dehydration amplifies anxiety, deepens low mood, and sharpens cravings, with a daily minimum of half your body weight in pounds, in ounces. For a Warrior: build genuine intensity (competition, mastery, vigorous exercise), because the pull toward stimulants is biological if it is not met. For a Sage: build calm (daily meditation, magnesium, nature, a structured routine, a low-caffeine life), because the Kernel names anxiety management as the central pillar. If focus was the reason, ask for an ADHD evaluation, since treating it removes the driver.

Turn to: Supp. Ch. 6 §4.1 (the two constitutions: the natural orientation and prevention priority). Supp. Ch. 20 §5 (the energy craving and the exercise-first response). Compendium Protocol 12 (Hydration Science), Protocol 19 (The Body in Motion), Protocol 03 (Sonic Architecture), Protocol 29 (The Peacetime Paradox).

Step 8 — If caffeine is bridging or swapping for something else

Two situations. If you are recovering from another substance and using caffeine to get through the fatigue, the Kernel is direct: caffeine does not deplete dopamine like methamphetamine does, so it is not "the same thing in miniature," but it stimulates the HPA axis on a system that may already be running hot (especially in a Met/Met "Worrier"), and heavy or late-day use undermines the sleep quality the repair depends on, so it plausibly lengthens the timeline indirectly. It says reaching for it in a hard stretch is understandable and not a moral failure, but it is not a neutral bridge. Managed, non-escalating caffeine can be a legitimate adrenergic tool for someone in the energy profile (the kratom guide's wording), but not as a substitute for the substance you left. If you are drifting the other way, from caffeine to alcohol to sleep, or to nicotine or cannabis, name the swap early, since the Kernel treats energy drinks and nicotine as entry points for the Warrior and caffeine as an amplifier for the Sage, and alcohol used for sleep suppresses REM.

Turn to: Supp. Ch. 32 §3 (the honest caveat: caffeine and alcohol as substitutes). Supp. Ch. 20 §5–6 (caffeine managed, for the energy profile). Supp. Ch. 6 (the entry-point warnings). Compendium Protocol 02 (Behavioral Addictions, with its "Mask Change" caution on addiction transfer).

Step 9 — Quarterly check and stewardship (Ongoing, Year 1 to Year 10)

The Kernel's quarterly self-check asks, among other things: have I noticed a substance producing a disproportionate pull, and have I been using caffeine, alcohol, or cannabis more than my baseline this quarter? For caffeine that means checking the ceiling, the noon cut-off, and whether sleep and anxiety are holding. Navigation after the first two months shifts from repair to stewardship. Section 4 shows how. Standing tools: Supp. Ch. 44 (The Transplant: redirect the devotion and daily structure the pattern required into a practice that grows). Supp. Ch. 6 (Stay Non-Addicted) and Compendium Protocol 02. Compendium Protocol 13 (The Pharma Trap). Supp. Ch. 38. Supp. Ch. 5, Step 6 (a weekly AI check-in profile, a reasoning tool and never a provider).

Turn to: The Kernel's quarterly self-check (the "substance relationship," "sleep quality," and "mood and anxiety baseline" rows). Supp. Ch. 6 (Stay Non-Addicted). Supp. Ch. 44 (The Transplant). Centurian Paper C-28 (The Recast).

4. How navigation changes at Years 3, 5, and 10

Shading shows how much attention each track needs. Darkest is primary focus, then active, then maintenance. White cells with gray text are standing safeguards: low effort, never dropped. Numbers in brackets refer to the shifts listed below the table. The Kernel gives no caffeine timeline, so this whole table is projection.

Track	Months 0–24 (foundation)	Year 3 (consolidate)	Year 5 (build out)	Year 10 (steward)
Safety and heart	Ceiling set; energy drinks out	BP and heart checked yearly [1]	Ceiling kept; no stacking	Noon cut-off; ceiling kept [7]
Biology and genome	CYP1A2, COMT; MTHFR check	Retest labs; step stack to upkeep [2]	Prescriber card: CYP1A2, COMT	Re-check with new meds and age
Timing and rituals	Noon cut-off; 3 PM cue broken	Ritual replaced; cues fainter	Triggers shift: deadlines, travel [5]	Same plan, rehearsed
Sleep and adenosine debt	Sleep debt repaid; light, timing	Sleep steady; no alarm needed [3]	Sleep habits held	Stress plan, sleep habits
Energy and calm architecture	Move for energy; meals, water	Own energy carries the day [4]	Calm practice daily	Movement, meals, still daily
Swap and substitute watch	Swap watch: alcohol, nicotine	Quarterly check: caffeine baseline	Quarterly check kept	Quarterly check kept
Purpose and Manning	Purpose: first seeds only	Sustainable work rhythm built	Manning: share what worked [6]	Mentor or run a group

The seven shifts

1. **Year 3, blood pressure and heart are checked yearly.** The Kernel flags elevated cardiovascular risk for slow CYP1A2 metabolizers at high doses (Supp. Ch. 6). By year 3 a routine yearly check of blood pressure is enough, tightened if you ever return to heavy use. This shift applies a Kernel rule at a timing chosen for this guide.
2. **Year 3, the stack moves from building to upkeep.** If you used supplements (magnesium, 5-MTHF), retest labs (Supp. Ch. 5 Step 1a) and step down toward maintenance. The Trinity chapter says supplements are taken until stable, then tapered and resumed if the system destabilizes; Main Book Ch. 2 treats MTHFR support as maintenance, not cure. Step down to maintenance, not to zero.
3. **Year 3, sleep is steady without props.** The Kernel says sleep is the primary repair intervention (Protocol 34 §34.5), and its own quarterly check asks whether you fall asleep within about 20 minutes and wake refreshed without an alarm. By year 3 that should be ordinary.
4. **Year 3, your own energy carries the day.** The Kernel's replacement for caffeine is exercise, meals, water, and the constitution's natural architecture

(Supp. Ch. 6 §4.1; Supp. Ch. 20 §5). By year 3 these are habits, not effort. This shift is extrapolation.

5. **Year 5, triggers shift to deadlines and travel.** The Kernel names a work deadline and the morning demand as when the energy craving fires (Supp. Ch. 20 §5), and shift work and travel as circadian disruptors (Supp. Ch. 6). By year 5, keep the noon cut-off and a plan for those weeks. This shift is extrapolation.
6. **Year 5, Manning.** This is the Trinity's third stage (Supp. Ch. 2). This is extrapolation: for caffeine it is modest and practical, sharing what worked with people whose sleep and anxiety are being eaten by the same pattern.
7. **Year 10, the noon cut-off and ceiling stay.** The Kernel's rule follows your enzyme, and your genotype does not change (Supp. Ch. 6). The quarterly check is the standing safeguard, and any new medication should be checked for interactions with caffeine, since CYP1A2 also clears some prescription drugs (Compendium §15.4).

What is sourced and what is extrapolated. *Shifts 3 and 7 rest directly on Kernel text. Shifts 1 and 2 apply Kernel rules at a timing chosen for this guide. Shifts 4, 5, and 6 are extrapolations. The Kernel gives no caffeine timeline.*

5. What outcome to expect

The short version

The Kernel has no caffeine case study, timeline, or percentages, so this section is qualitative and does not make up numbers. If you do the route above, with the ceiling and cut-off set, the genes known, sleep repaired, and energy and calm built in other ways, this is what the Kernel's material supports:

- **Sleep improves first.** The Kernel treats it as the primary repair intervention, and late caffeine is the most direct thing you control.
- **Anxiety and thought "noise" ease,** especially for a slow-COMT profile, because caffeine raises excitatory activity and lowers GABA function.
- **The heart risk falls** if you were a slow metabolizer using large energy drinks.
- **By years 3 to 5 the work is maintenance and building. By year 10 you are a steward,** carrying two standing safeguards: the noon cut-off and the quarterly check.

The Kernel does not promise that caffeine has to be zero. It says the right amount depends on your constitution, and that for some people, the right amount is none.

Milestones, in order

Milestone	When	Expected outcome	Source	Basis
Heart and safety checked	Day 0–7	Form and amount known. Heart symptoms and pregnancy raised with a clinician. No stacking.	Supp. Ch. 6	Clinical fact

Milestone	When	Expected outcome	Source	Basis
Withdrawal window	Days 1–7 (about)	Headache, fatigue, low mood, and poor concentration are typical, and fade within about a week. Gradual cutting is gentler.	Not in the Kernel	General medical knowledge
Ceiling and cut-off in place	Weeks 1–2	Noon cut-off. Ceiling set (Kernel: 100–150 mg/day for a slow CYP1A2). No very large energy drinks. Sleep and meals regular.	Supp. Ch. 6	Kernel projection
Sleep improves	Weeks 2–8	Sleep debt repaid. Falling asleep and waking without an alarm become normal. Anxiety and thought noise ease.	Protocol 34 §34.5	Kernel projection
Own energy	Weeks 4–12	Exercise, meals, water, and the constitution's architecture carry the day without a stimulant.	Supp. Ch. 6 §4.1; Ch. 20 §5	Kernel projection
Steward phase	Years 3–10	Noon cut-off and ceiling kept. Quarterly check. Annual blood-pressure check.	Section 4 of this guide	Extrapolated

What changes the outcome

- **Late-day caffeine,** especially in a slow CYP1A2 metabolizer. It steals the sleep that repair needs.
- **Very large or stacked energy drinks** and caffeine powders or pills. The Kernel names the heart risk.
- **Ignoring COMT.** A Sage is amplified by caffeine, and a Warrior may be drawn from coffee to harder stimulants.
- **Using caffeine to power through fatigue** during recovery from another substance.
- **Swapping to alcohol for sleep,** or to nicotine or cannabis, and not naming it.
- **Synthetic folic acid** in energy drink blends, for MTHFR variants.
- **Skipping meals and water,** which the Kernel says amplify anxiety and cravings.
- **Treating a bad night as a reason to double up.** The debt is collected later, with interest.

How to read the evidence behind the outcomes

The Main Book marks claims with an Evidence Tier: Established, Probable, Emerging, Author/System Experience, and Controversial or Mixed. This guide's reading: caffeine's adenosine blockade, its effect on sleep, its recognized withdrawal, and the higher cardiovascular risk in slow CYP1A2 metabolizers at high intake are supported in the wider literature, as I understand it, though the size of the CYP1A2 effect varies. The Kernel's COMT-and-caffeine advice is its own inference from the Warrior and Sage framework and sits lower on the scale. Some of its Bartender's Friend coffee claims look shaky: coffee is not a significant source of tyrosine, caffeine is cleared by CYP1A2 and not CYP2D6 (so a CYP2D6 "rapid metabolizer" is not a caffeine rapid metabolizer), and I could not confirm that cold brew clears more slowly than espresso. The numbers in the Kernel's CYP1A2 row (a 100 to 150 mg ceiling, none after noon) are its rules of thumb, not trial results. Bring all of this to your clinician as questions, not as facts.

Take these to your clinician

- Every source of caffeine you use, with rough amounts and times, including energy drinks, pre-workout, pills, and powders.
- Any palpitations, chest pain, high blood pressure, anxiety, or trouble sleeping, and any other stimulants or stimulant medications you take.
- Your CYP1A2 and COMT results if you have them, and whether pregnancy or a heart condition applies.
- A question about whether your focus or fatigue needs its own evaluation (ADHD, sleep, iron, thyroid, or depression), since caffeine may have been covering it.